

## (1) Submission ID#1539536

*Neisseria lactamica* and protection against meningococcal colonisation

---

### Author(s)

Adam Dale, PhD, BSc, MBChB, MRCP, FRCPATH

Clinical Lecturer

University of Southampton

Jay Laver, BSc PhD

Senior Research Fellow

University of Southampton

Diane F. Gbesemete, BM, MRCPCH, PGDipID

Clinical Research Fellow

University of Southampton

Anastasia Theodosiou, MBBChir

Research Training Fellow

University of Southampton

Andrew Gorringe, PhD

Scientific Leader

UK Health Security Agency

Saul Faust, PhD

Professor of Paediatric Immunology and Infection

University of Southampton

Robert Read, BMedSci(physiology), MBChB, MRCP, MD, FRCP(London), FRCP(Glasgow)

Professor

University of Southampton

### Background

Unlike the potent glycoconjugate vaccines, subcapsular antigen-based vaccines targeting serogroup B *N. meningitidis* (Nm) strains do not protect against pharyngeal colonisation and are therefore unlikely to induce herd immunity. For optimal public health control of serogroup B disease, alternative strategies will be required. Natural, and experimentally induced colonisation with *Neisseria lactamica* (Nlac) provides broad spectrum protection against Nm colonisation. Nlac formulated as an intranasally-administered Live Bacterial Product (LBP) could feasibly be utilised to limit Nm carriage. Here, we demonstrate that: (1) the protective effect of Nlac on Nm may be due to induction of cross-reactive adaptive immunity, (2) anti-Nm adaptive immunity

generated following Nlac colonisation can be enhanced through genetic modification of Nlac.

#### Aim/Methods

We conducted two controlled human infection experiments using either wild type (WT) or genetically-modified Nlac (GM-Nlac) expressing the meningococcal vaccine antigen, NadA.

#### Results

Colonisation with WT Nlac induced anti-Nlac- and anti-Nmen-specific IgG, plasma cells (BPLAS) and memory B cells (BMEM). B cell dynamics suggested that Nlac colonisation recalled pre-existing Nmen-cross-reactive IgG BMEM. Nlac colonisation density correlated inversely with the magnitude of anti-Nlac immunoglobulin titres and BPLAS responses, suggesting a role for these responses in bioburden control. Colonisation with GM-Nlac was safe and generated anti-NadA IgG, BPLAS, and BMEM. In contrast to WT Nlac, GM-Nlac induced serum bactericidal antibodies against serogroup B strain 5/99.

#### Conclusions

Nasal inoculation with both WT Nlac and GM-Nlac is safe and protects against Nmen colonisation. Current evidence suggests this effect is driven by generation of cross-reactive adaptive immunity. The observation that GM-Nlac induced anti-NadA adaptive immune responses, including generation of SBA titres, justifies further studies to assess the impact of GM-Nlac on protection against Nmen colonisation and disease. To that end, we will discuss current progress in the development of GM-Nlac expressing multiple Nmen antigens (including NadA and fHBP) suitable for use in controlled human infection, experimental medicine and as a potential LBP.