

(1) Submission ID#1527817

Structure of anti-gonococcal lipooligosaccharide antibody bound to mimetic peptide vaccine informs therapeutic and vaccine strategies

Author(s)

Peter T. Beernink, Ph.D.

Professor

University of California, San Francisco

Cristina Di Carluccio, Ph.D.

Postdoctoral Scientist

University of Naples Federico II

Roberta Marchetti, Ph.D.

Professor

University of Naples Federico II

Benjamin Beernink, Undergraduate Student

Student Intern

University of Callifornia, San Francisco

Sunita Gulati, DSc

Professor

Division of Infectious Diseases and Immunology, University of Massachusetts Chan Medical School,
Worcester MA, USA

Jutamas Shaughnessy, MD, PhD

Associate Professor

Division of Infectious Diseases and Immunology, University of Massachusetts Chan Medical School,
Worcester MA, USA

Peter Rice, MD

Professor

Division of Infectious Diseases and Immunology, University of Massachusetts Chan Medical School,
Worcester MA, USA

Sanjay Ram, MBBS

Professor; Department of Medicine; Division of Infectious Diseases and Immunology

University of Massachusetts Chan Medical School

Alba Silipo, Ph.D.

Professor

University of Naples Federico II

Background

Neisseria gonorrhoeae (Ng) is the etiologic agent of gonorrhea, which causes local genital infection and disseminated disease. Mouse monoclonal antibody (mAb) 2C7 recognizes a widely expressed glycan epitope of Ng lipooligosaccharide (LOS), called the 2C7 LOS epitope, and elicits complement-dependent bactericidal activity. Previously, we identified a cyclic peptide called CP2 that mimicked the 2C7 LOS epitope. Immunization of mice with CP2 configured as a tetrameric multi-antigen peptide, TMCP2, attenuated Ng vaginal colonization.

Aim/Methods

We used biophysical approaches to characterize the interactions between the therapeutic mAb candidate 2C7 and the multi-antigenic mimetic peptide vaccine candidate TMCP2. To this end, we employed small-angle X-ray scattering, native mass spectrometry, NMR spectroscopy, X-ray crystallography and molecular dynamics (MD) simulations to characterize complexes of mAb and Fab 2C7 with tetrameric and monomeric peptides TMCP2 and CP2.

Results

Small-angle X-ray scattering experiments with Fab 2C7 identified changes in the radius of gyration and shape upon forming complexes with CP2 and TMCP2. Native mass spectrometry with mAb and Fab 2C7 displayed different oligomeric structures bound to tetrameric TMCP2. Saturation transfer difference and transferred nuclear Overhauser effect NMR spectroscopy showed that the chemical environments of primarily hydrophobic CP2 residues changed with mAb 2C7 binding. The high-resolution crystal structure of Fab 2C7 in a complex with CP2 revealed that the peptide formed a beta-hairpin and confirmed that the Fab bound via hydrophobic interactions. A second crystal structure of Fab 2C7 alone showed that there were no substantial conformational changes in the Fab upon peptide binding. Finally, NMR studies, crystallographic thermal factors, and molecular dynamics simulations revealed the dynamic conformational behavior of CP2 when bound to mAb/Fab 2C7.

Conclusions

Using a combination of biophysical approaches, we mapped the CP2/TMCP2 epitope and characterized the interactions between mAb 2C7 and CP2. These studies, in conjunction with MD simulations, identified the bioactive conformations of CP2 when bound to Fab 2C7. Collectively, our studies will guide strategies for humanizing mAb 2C7 for use as a therapeutic antibody against gonococcal infection, and for optimizing CP2 as a gonococcal vaccine candidate.

Uploaded File(s)

Supplemental Document Upload

[Structure of mimetic peptide CP2 \(red\) bound to Fab fragment of therapeutic mAb 2C7 \(green and blue\).](#)

