

(1) Submission ID#1526711

MenB-4C vaccination elicited IgG antibodies recognize meningococcal and cross react with gonococcal lipooligosaccharides (LOS)

Author(s)

Yih-Ling Tzeng, Ph.D.

Research Associate Professor

Emory University School of Medicine

Soma Sannigrahi, n/a

Senior Research Specialist

Emory University School of Medicine

Ray Borrow, Ph.D.

Professor

Meningococcal Reference Unit, UK Health Security Agency

Nadine Rouphael, M.D.

Professor

Emory University School of Medicine

Varun Phadke, M.D.

Assistant Professor

Emory University School of Medicine

David Stephens, M.D.

Professor

Emory University School of Medicine

Background

Antigens in the outer membrane vesicles (OMVs) of the *Neisseria meningitidis* (Nm) group B-directed vaccine MenB-4C (Bexsero) may protect against infections with *N. gonorrhoeae* (Ng). The immunologic basis for this protection remains unclear. The lipooligosaccharide (LOS) of vaccine strain NZ98/254 is a mixture of L1 and L3 structures with an inner core 3-PEA-HepII modification. Previous work has shown that the OMV-based vaccines generate serum antibodies to OMV proteins and LOS .

Aim/Methods

The aim of this study was to characterize the human serum IgG antibodies against Nm and Ng LOSs elicited by MenB-4C. Ten paired sera obtained pre- and post-immunization (one month after a third vaccine dose, Vaccine 2015;33(29):3322-30) were used in Western blots to probe Nm and Ng whole cell lysates after

removal of proteins by protease K. Three post MenB-4C sera were then used to probe the structurally defined meningococcal LOS immunotypes (L1-L12) and LOSs of Nm and Ng strains and mutants.

Results

Post (but not pre-immune) MenB-4C sera recognized both Nm and Ng LOS species, accounting for ~5-9% of total IgG response to gonococcal OMV antigens. Nm and Ng LOSs were broadly recognized by post-immunization IgG antibodies but with individual variability in structural specificity. Terminal sialylation of the α -chains had minimal impact on recognition. A maltose (Glc-Glc; L5, L10 and L11), instead of lactose (Gal-Glc), linkage to HepI significantly diminished or eliminated reactivity. Deep truncation of LOS, specifically the *rfaK* mutant without glycosylation of α -, β - or γ -chains, was not recognized by any post vaccine sera. The 3-PEA-HepII could be part of the conformational epitope; serum 7v5 IgG antibodies predominantly recognized the L1 (Gal-Gal-Glc-HepI) α -chain with a 3-PEA-HepII β -chain. Replacing with a 3-Glc at HepII blocked antibody recognition. Serum 19v5 recognized inner core LOSs with truncated α -chain structures (Gal-Glc-HepI, L8) and the 3-Glc-HepII also interferes with antibody binding. LOS antibodies in serum 17v5 recognized an intact α -chain, either L3 lactoneotetraose (LNT) or L1, and contributed to serum bactericidal activity against gonococci.

Conclusions

MenB-4C vaccination elicited serum IgG antibodies to conformational epitopes involving HepI and HepII glycosylated LOS structures shared between Nm and Ng that may be a bactericidal target.