

(1) Submission ID#1539422

Development of a gonorrhoea pharyngeal controlled human infection model for the assessment of novel prevention and therapeutic strategies

Author(s)

Georgina Pollock, PhD

Research Fellow

Department of Infectious Diseases, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia

Eloise Williams, MBBS, BMedSci, MPH, FRACP, FRCPA

Clinical Microbiologist and Infectious Diseases Physician

1. Department of Infectious Diseases, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia; 2. Victorian Infectious Diseases Reference Laboratory, Royal Melbourne Hospital at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia

Soo Jen Low, PhD

Research Officer

Department of Infectious Diseases, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia

Shivani Pasricha, PhD

Senior Research Officer

Department of Infectious Diseases, The University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Australia

Efrosinia O. Krejany, BSc, MSc

Clinical Research Operations Manager

Department of Infectious Diseases, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia

Mona Taouk, BBiomed [Hons]

PhD student

Department of Infectious Diseases, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia

Julia Marshall, MBBS, MPH, FRACGP, DipObs, PhD

Clinician and Post-Doctoral Scientist

Department of Infectious Diseases, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia

Deborah Williamson, BSc MBChB MRCP FRCPA PhD

Director of the Victorian Infectious Diseases Reference Laboratory; Professor of Microbiology

1. Victorian Infectious Diseases Reference Laboratory, Royal Melbourne Hospital at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia; 2. Department of Infectious Diseases, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia 3. The Walter and Eliza Hall Institute of Medical Research, Melbourne, Victoria, Australia

James McCarthy, MBBS, FRACP, PhD

Director of the Victorian Infectious Diseases Service; Professor of Medicine

1. Victorian Infectious Diseases Service, Royal Melbourne Hospital, Melbourne, Victoria, Australia; 2. Department of Infectious Diseases, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia; 3. The Walter and Eliza Hall Institute of Medical Research, Melbourne, Victoria, Australia

Background

Infection with *Neisseria gonorrhoeae* is a major public health problem characterised by increasing global incidence and antimicrobial resistance. Accordingly, novel and innovative gonorrhoea prevention and therapeutic strategies are urgently required. As *N. gonorrhoeae* is a human-restricted pathogen, preclinical animal infection models for assessment of novel interventions have important limitations. A gonorrhoea urethritis controlled human infection model (CHIM) has led to significant progress in understanding gonorrhoea urethritis pathogenesis. Increasing evidence suggests that pharyngeal infection plays a key role in gonorrhoea transmission and antimicrobial resistance, however there is a paucity of data describing *N. gonorrhoeae* pathogenesis and host responses to infection of the pharynx. Here, we describe progress towards establishing a first-in-field pharyngeal gonorrhoea CHIM.

Aim/Methods

First, genomic analysis will be used to select a panel of contemporary, globally-relevant *N. gonorrhoeae* strains optimised for patient safety and vaccine antigen coverage from a panel of 5,911 *N. gonorrhoeae* isolates. This panel will undergo detailed phenotypic characterisation including antimicrobial susceptibility testing and in vitro infection assays using human pharyngeal epithelial cell lines. In parallel, development and validation of standard operating procedures for preparation and storage of the challenge strains will be performed in accordance with good manufacturing practice (GMP), aimed at obtaining regulatory approval for challenge strain production. An ethical and locally-appropriate study design for a pharyngeal gonorrhoea CHIM will be then be developed, informed by a program of community engagement and consultation with key stakeholders. Finally, an observational, dose-escalation inpatient pharyngeal gonorrhoea CHIM trial will be undertaken in carefully-screened healthy adult male volunteers.

Results

The requirements for a contemporary pharyngeal gonorrhoea CHIM will be described, including i) the rational selection of a contemporary, globally-relevant *N. gonorrhoeae* strain; ii) development of a GMP-grade cell bank that produces well-characterised single-dose vials of the challenge strain and iii) design of an ethical and locally-appropriate clinical trial protocol.

Conclusions

Development of a contemporary gonorrhoea pharyngeal CHIM may potentially serve as a powerful platform for assessment of novel gonorrhoea prevention and therapeutic strategies.